

Morphological State of Kurashs with Evaluation of Sports Specialization

*Keldiyorov Shuxrat Ergashovich, Yusupov G'ayrat Abdullaevich,
Bobomurodov Feruz Ismoiljonovich*
Republic of Uzbekistan, Chirchik, Sportchilar Street, 19

ABSTRACT

Categories of somatotypes of athletes specializing in various types of kurash are established by the methods of anthropometry and by Hit-Carter method. Results of anthropometric parameters showed that somatic state of athletes engaged in various types of kurash are differed. Similarity was found in constitution of kurashs specializing in weightlifting, freestyle kurash and national sport kurash, and endomesomorphic somatotype was determined in 67-83% of athletes, that calculating by points is expressed as 3.3:6:2. Ecto-mesomorphic somatotype dominated among judoists and sambo kurashs and components of body mass were presented as 3,3: 5,4: 2,2 and 2.9: 5.4: 2.3 respectively. Dynamic observation showed that athletes with balanced-mesomorphic somatotype possess wide adaptation ability, providing an effective rearrangement of morphotype under the influence of training modes of different orientation.

ARTICLE INFO

Article history:

Received 11 Sep 2024

Received in revised form

15 Sep 2024

Accepted 24 Oct 2024

Keywords: relevance, variability, wrestling, integral, characters, transverse, empirical, respectively, length, somatotype, balanced, comparative, fact, category.

Hosting by Innovatus Publishing Co. All rights reserved. © 2024

The relevance of research: Parameters of body type occupy an important place among factors influencing sport activity. Variability of body size depend on age, profession and they could be of selective significance in sport [(1,6,8]. Somatotype as an integral characteristic of the athlete morphological organization, is a marker of motor and some functional capabilities of the organism. Genetic and environmental factors influence somatotype formation [2. 9,]. In this connection one of the most important environmental factors is the purposeful sports activities forming adaptive somatotype of an athlete of specific specialization [3, 6, 7, 12]. Analysis of morphological characters should be conducted in various aspects: 1. analysis of morphologic characters during process of individual development taking into consideration critical and sensitive periods of development; 2. tracing the formation of a certain body type under the influence of training, characteristic for a certain sport specialization, or the use of morphological features for the purposes of athletic selection and orientation. Condition of physical development of athletes is characterized by a number of anthropometric signs. There are single data on a possibility of significant changes not only in the component composition of the body, but also in the type of constitution, possibility of goal-directed changes of junior constitution as a result of practicing a certain sport [4].

In this regard, the task was set: to find out morphological differences among representatives of the different specialization, in particular in various kinds of wrestling, to study somatotypes variations, changes in the composition of the body mass of kurashs, depending on the different orientation of the training regimes in the course of long-term dynamic observations.

The following research methods were used:

1. **Anthropometric studies.** Estimation of physical development of athletes was carried out by the methods of anthropometry. Anthropometric studies were conducted according to the methods described in the manual of E.G. Martirosov (1982) [5] and using standard tools. The athletes under examination

represented an ethnically homogeneous group, as medical and biological approach to the problem of sport orientation and selection assumes consideration the individual characteristics of a person, formed on biological prerequisites, having a genetic basis. Athletes specialized in power sports: weightlifting (35 athletes) and 5 types of kurash (153 athletes). In total 288 athletes were under examination. Total sizes, as well as partial: longitudinal, transverse, girth of the trunk, head, neck, segments of the upper and lower extremities were measured. The thickness of the skin-fat folds was measured in 4 parts of the body: under the scapula, in the upper iliac region, on the posterior surface of the shin and shoulder. Main anthropometric parameters and mean square deviation were calculated.

2. Type of constitution, detected by the Hit-Karter method, 1989 [11, Carter J.L.], is one of universal methods of somatotypology as it is characterized by common quantitative approaches for the determination of variants of constitution in people of all nationalities and races, of any age and sex. The Hit-Carter scheme is universal also because it is based on 7 anthropometric features and covers the entire empirical set from minimum to maximum values and the principle of approach to the construction of tables. It means that results obtained in examination of athletes of various specialization, are comparable as well as data obtained in athletes at the period rehabilitation and during active training. Besides this scheme has become very popular in foreign studies, i.e. there is the opportunity to compare own research with research conducted in other laboratories.

Standard measurements were originally conducted as detection of length of the body, diameter of the distal part of the shoulder, thigh, shoulder circumference in the stressed state, shin circumference, and the caliper measurements of the skin-fat folds in 4 regions of the body were also carried out.

Somatotype was diagnosed on the basis of three components: I IF is the fat component of endomorphism; II M - muscular component - mesomorph; III -Ectomorphy - P/L -weight-growth index. In the process of dynamic observations after 2 years of occupation by the selected sport, detection of the component composition of the body mass with the subsequent diagnosis of the somatotype was performed.

Results and discussion. Analysis of obtained results showed that in athletes engaged in strength sport, freestyle wrestling, kurash, Greco-Roman wrestling, the length of the body is approximately the same: $175.72 + 0.62$ (freestyle wrestling), $175.30 + 2.52$ (kurash), $175.44 + 1.08$, (Greco-Roman), $175.44 + 1.08$ (weightlifters), respectively, exception was presented by judoists (Table 1). However in spite the similar body length, body mass varied in all kinds of sport. the mean square deviation ranged from 7.99 to 11.04.

High variability in chest circumference was established for freestyle wrestlers, whose standard deviation was 9.89, for sambo wrestlers -6.54 and for weightlifters -6.86.

A similarity in the proportions of the body is established. The most stable were partial long lengths: the length of the shoulder, the forearm, the shin, the circumference of the head, neck, shin, shoulder width. Variability is revealed in the dimensional characteristics of the wrists –the range in the mean square deviations was from 1.20 to 9.90.

Evaluation of the dimensions of fat folds in the compared groups showed uniformity in the quantitative distribution of the subcutaneous fat layer, except for the thickness of the fat fold under the scapula. In weightlifters the thickness of the fold was $9.05 + 2.39$, for kurashes this value was $7.96 + 2.33$. Intergroup variability is found in the parameters of the abdomen. This parameter in weightlifters amounted to $81.80 + 3.9$ cm, for freestyle wrestlers $-79,40 \pm 0,71$, for sambo wrestlers $-83,20 \pm 1,43$, for Greco-Roman wrestlers $-78,00 \pm 0,42$, for kurashes $-75,40 \pm 1,10$. Differences in length of the hip and width of the pelvis were also revealed. So weightlifters are characterized by a larger pelvic width than kurashes, which is associated with the specificity of training, in particular, with the development of a "low sub" under the bar. Analysis of morphometric characteristics of kurashes in 5 kinds of kurash was carried out. Differences in mean square deviation values are revealed. Thus, the circumferential dimensions of the abdomen, shoulder, hand, hip in specialization types under consideration repeat properties of intergroups distribution.

The range of variation of features is characteristic for indices of length of the forearm, shin, neck circumference, diameters of the middle chest, apical joint, thickness of the fat fold on the posterior surface of the shoulder (Table 1). Probably these signs are more affected by the specificity of sports

specialization, the character of the training regimes, choice of a specific arsenal of attacking technical actions.

Analysis of somatotypes categories demonstrated domination of ecto-mezomorphic type among judoists and sambo wrestlers (33.4-41.7%). An equal distribution of two somatotypes is observed in classic wrestlers: ecto-mesomorphism and balanced mesomorphy occurred in 50% and 41,76% of cases, respectively. Calculation of the main somatotype components by points showed dependence on the specifics of the specialization. Thus endo- and mezomorphic components achieve the maximal development in free style wrestlers (3,3: 6,0:2,1), less values were detected in kurashs engaged in kurash and sambo: 3,3: 5,4: 2,2; 2,9: 5,4: 2,3 respectively. Contrary to the popular opinion that intensive sport exercises contribute to maximal fat decrease in an organism, there is convincing evidence that for representatives of some kinds of sports, in particular, for kurashs and weightlifters, the amount of fat can be quite large, since fat is the main energy potential. Higher values of ectomophic component are typical for classic wrestlers, in which the average values of the somatotype in points are 2.8: 4.6: 2.7. Obtained results indicate that degree of similarity and differences in values of anthropometric parameters depend on the nature of the sign. .

Comparative analysis of kurashs somatotyping in 2 years allowed to establish the maximal differences in degree of metamorphic component expression. Minor differences were identified in indices of endomorphism. Registration of category of somatotypes revealed the following:

at the initial stage category of endomorphic type of kurashs amounted to 65.2%, and in 2 years this value achieved 82.6% at the expense of transition of balanced mezomorphic type into endo-mezomorphic type.

Somatic polymorphism is observed in kurashs who have even insignificant differences at the level of athletic skill, and changes in somatotypes are noted in athletes whose adaptive capabilities provide an effective morphotypic restructuring, more optimal for the selected sport. Such differences among athletes who specialize in different types of kurash are not only determined by technical and tactical expediency, but they are also connected with the morphological prerequisites of development.

The revealed facts once again confirm the influence of exogenous factors on the program of individual ontogenetic development. In this case, the influence of the exogenous factor is realized through the improvement of motor, speed-strength qualities in the direction necessary for a particular sport. The orientation of the training process, the change in the technique of motion even within the same specialization, leads to a sharp alterations in the component composition of the body, which ultimately leads to a change in the category of the somatotype.

Conclusion

1. Somatotype polymorphism among athletes specializing in 5 types of kurash was established.
2. Endo-mesomorph category dominated in athletes engaged in free style kurash, kurash and weightlifters. Ecto-mesomorphic type was to a large extent represented by Sambo wrestlers and judoists. Endo-mesomorphic and balanced-mesomorphic types were revealed in free style kurashs
3. Athletes with balanced - mesomorphic types have wide adaptive capabilities that ensure the effective restructuring of the morphotype under the influence of training regimes of different directions.
4. Somatic states are differed in athletes engaged in various kinds of kurash and they can be considered as separated classes.

References

1. Dorohov R.N. Basis and prospects of age somatotyping. Theory and practice of physical culture. 2000. N0. P.10;
2. Cogan B.IU. Hereditary and environmental conditioning of body elements. News in sport medical anthropology. M., 1990, issue 1. P. 83-84.
3. Martirosov E.G. Systemic organization of somatic state of athletes and classification of sport specializations. Morphologic problems of sport selection. Collection of research papers. Moscow. 1989. P.5-30.

4. Martirosov E.G. Methods of research in sports anthropology. Moscow, FiS, 1982.
5. Nikityuk B.A. Constitutional aspects of anthropology. Integrative biosocial anthropology" Moscow, 1996. - 220 p.
6. Pavlova O.I. Present day training of young athletes. Moscow, publishing house "Theory and practice of physical culture", 2004 - P. 32-34.
7. Panasyuk T.V. , Tambovtzeva R.V. Constitutional features of the physical performance of adolescents. VII International Scientific Congress "Modern Olympic Sports and Sport for All", Volume II, Moscow, 2003. - 135 p.
8. Rogozkin V.A. et al. Genetic markers of a person's physical working capacity. // Theory and practice of physical culture. - 2000. N. 12. P. 34 - 36.
9. Rogozkin V.A. Decoding of the human genome and sport. Theory and practice of physical culture, 2006, №6, - P. 60-63.
10. Carter J.L. – “Somatotype characteristics of champion athletes”, Rep, Antrop-Cong. ed. A. Hrdica, Praha, Humpolec. 1969.
11. Timility D. Protocols for Physiological assessment of male and female soccer players. In Physiological tests for Elite Athletes, Human Kinetics, 2000, 17, 356-362.